arXiv:0712.2236v2 [gr-qc] 6 Mar 2008

Cosmic string loops: large and small, but not tiny

Vitaly Vanchurin*

Arnold-Sommerfeld-Center for Theoretical Physics, Department für Physik,
Ludwig-Maximilians-Universität München, Theresienstr. 37, D-80333, Munich, Germany

We develop an analytical model to study the production spectrum of loops in the cosmic string network. In the scaling regime, we find two different scales corresponding to large (one order below horizon) and small (few orders below horizon) loops. The very small (tiny) loops at the gravitational back reaction scale are absent, and thus, our model has no ultra-violet divergences. We calculate the spectrum of loops and derive analytical expressions for the positions and magnitudes of the small and large scale peaks. The small loops are produced by large bursts of similar loops moving with very high velocities in the same direction. We describe the shape of large loops, which would usually consist of few kinks and few cusps per oscillation cycle. We also argue that the typical size of large loops is set by the correlation length, which does not depend on the intercommutation probability p, while the interstring distance scales as $p^{1/3}$.

I. INTRODUCTION

Many cosmological models lead to the production of cosmic strings as a result of the phase transition [1], or cosmic superstrings at the end of brane inflation [2, 3, 4]. Very early analytical [1, 5, 6] and numerical [7, 8, 9] investigations agreed that the evolution of infinite strings exhibits scaling. An unexpected result came from the analysis of string loops by means of numerical simulations, when the two groups [8, 9] independently found that the loops do not scale, and are mostly produced at the smallest resolution scale of the simulation.

To tackle this problem an exact simulation for the cosmic string network in flat space-time [10] was developed. The precision of the simulation was only set by unavoidable computer arithmetic uncertainty, which is many orders of magnitude smaller than the scale of any structure in the network. Using this code, it was shown that the production of small loops is always accompanied by the production of somewhat larger loops [11]. Later, it was conjectured [12] that the production of small loops is a transient phenomena and in the long run only the large loops are produced at the scales of the interstring distance $\sim 0.1t$.

The two other groups [13, 14] have recently performed the numerical analysis with higher resolution versions of the simulation codes previously used [8, 9] and were able to resolve the small loops. Reference [13] found some evidence for a scaling distribution of loops, but their size is apparently much smaller than in Refs. [11, 12] and Reference [14] found a distribution which diverges at small sizes. We believe that the dynamical range of all three simulations [9, 12, 14] is not sufficient, to study the true scaling behavior.

In an attempt to settle the issue, in this article we develop an analytical model of cosmic strings and compare it with another recently proposed model of Ref. [16, 17]. We show that in the final scaling regime both types of loops (small and large) are present, and we do not see any evidence of the tiny loops at the gravitational back reaction scale.

The paper is organized as follows. In the next section we analyze the production of large and medium-sized loops. In the third section we study the production of small loops in the vicinity of cusps. We show that the dominant scale of small loops is only few orders smaller than horizon, and the production of loops at the gravitational back reaction scale is strongly suppressed. In the forth section we calculate the spectrum of loops in three different regimes with intercommutation probability also taken into account. In the fifth section we derive analytical expressions for the positions and magnitudes of the small and large scale peaks in the power law approximation of the correlation functions. In the last section we briefly discuss the effects of cross-correlations between opposite-moving waves and compare our results with those obtained in Ref. [16, 17].

II. LARGE SCALES

The evolution of cosmic strings can be captured by right ($\mathbf{a}(\sigma)$) and left ($\mathbf{b}(\sigma)$) moving waves

$$\mathbf{x}(\sigma,t)=\frac{1}{2}(\mathbf{a}(\sigma-t)+\mathbf{b}(\sigma+t))$$

where $\mathbf{x}(\sigma,t)$ describes the shape of a given string. When a self-intersection takes place, the following condition must be satisfied

$$(\mathbf{a}(\sigma+\Delta-t)-\mathbf{a}(\sigma-t))+(\mathbf{b}(\sigma+\Delta+t)-\mathbf{b}(\sigma+t))=0$$

where Δ is the size of a loop that could be formed, and t is the time of the intersection.

For convenience, we define

$$\mathbf{A}(\sigma)\equiv\int_{\sigma}^{\sigma+\Delta}\mathbf{a}'(x)dx=\mathbf{a}(\sigma+\Delta)-\mathbf{a}(\sigma)$$

---

*Electronic address: vitaly@cosmos.phy.tufts.edu

and

$$\mathbf{B}(\sigma) \equiv \int_{\sigma}^{\sigma+\Delta} \mathbf{b}'(x)dx = \mathbf{b}(\sigma+\Delta) - \mathbf{b}(\sigma)$$

The condition of self-intersection can be replaced by an inequality

$$\mathbf{A}(\sigma - t) + \mathbf{B}(\sigma + t) < \epsilon,$$

which must be satisfied for some σ and t in order for a loop of a size from Δ to $\Delta + \epsilon$ to have a chance to be formed. In the limit of small ϵ, the self-intersection takes place with probability approaching to one. If the above inequality holds for an arbitrary small ϵ, then we say that a loop is formed with probability p, where p is determined from the underlying theory.

In general, the vectors $\mathbf{A}(\sigma)$ and $-\mathbf{B}(\sigma)$ lie within a volume V of a sphere with radius Δ. However, for the extreme scales corresponding to very large (Brownian) or very small Δ, the vectors are confined to a somewhat smaller sphere or to a spherical shell respectively. In this article we model the production of loops as a random process of finding $\mathbf{A}(\sigma)$ and $-\mathbf{B}(\sigma)$ such that their difference is smaller than ϵ. The intuition tells us, that it must be easier to find the right counter-partner if the volume V is smaller. However, this is not the whole story and other factors must be taken into account in order to avoid divergences at small scales.

If the vectors $\mathbf{A}(\sigma)$ and $-\mathbf{B}(\sigma)$ span the entire sphere of radius Δ, then we call loops formed at that scale, the medium-sized loops and

$$V_{medium} = \frac{4}{3}\pi\Delta^3.$$

Large loops, are the loops formed from the vectors $\mathbf{A}(\sigma)$ and $-\mathbf{B}(\sigma)$, which are confined to a sphere of radius $\propto \xi\sqrt{\frac{\Delta}{\xi}}$, where ξ is the correlation length along the string and

$$V_{large} = \frac{4}{3}\pi\Delta^{\frac{3}{2}}\xi^{\frac{3}{2}}.$$

Some of the large loops do not inter-commute with infinite strings throughout their evolution, when other quickly rejoin back to the network. As we are only interested in the loops that are not affected by the infinite strings, it is reasonable to expect a sharp cut-off at

$$\alpha_l \sim \pi\xi, \tag{1}$$

for the interstring distance of order ξ and intercommutation probability $p = 1$.

The very large loops are not stable due to their Brownian shape, regardless of p. After few cycles of oscillations the loops fragment into smaller loops, which can only decay further by gravitational radiation. Therefore, the large loops in the final spectrum (after fragmentation) are likely to consist of relatively straight segments with at most few kinks (~ 4) and cusps (~ 2 per oscillation cycle), which is in agreement with numerical simulations of Refs. [11, 12].

III. SMALL SCALES

For small loops, the volume occupied by vectors $\mathbf{A}(\sigma)$ and $-\mathbf{B}(\sigma)$ is approximated by a spherical shell of radius Δ and thickness δ with

$$V_{small} = 4\pi\lambda_1^2\delta,$$

where

$$\begin{aligned}\lambda_1 &\equiv \langle\mathbf{A}\rangle_{rms}\\ &= \langle\int_{\sigma}^{\sigma+\Delta}\mathbf{a}'(x)dx\rangle_{rms} =\\ &= \langle\mathbf{a}(\sigma+\Delta) - \mathbf{a}(\sigma)\rangle_{rms}\end{aligned}$$

and

$$\delta \equiv \sqrt{\langle\mathbf{A}^2\rangle - \langle\|\mathbf{A}\|\rangle^2}.$$

Thus, nearly all of the small loops are emitted with very high velocities by combining nearly straight segments of **a**- and **b**-waves pointing in opposite directions. In such regions the following condition is likely to be satisfied satisfied

$$\mathbf{a}'(\sigma) = -\mathbf{b}'(\sigma),$$

which indicates the presence of a cusp.

Without loss of generality, we consider **a**- and **b**-segments of the string such that

$$\begin{aligned}\mathbf{a}'(0) &= -\mathbf{b}'(0)\\ \mathbf{a}(0) &= 0\\ \mathbf{b}(0) &= 0\end{aligned}$$

and expand it in powers of σ

$$\begin{aligned}\mathbf{a}(\sigma) &= \sigma\,\mathbf{a}'(0) + \frac{\sigma^2}{2}\,\mathbf{a}''(0) + \frac{\sigma^3}{6}\,\mathbf{a}'''(0) + ...\\ \mathbf{b}(\sigma) &= \sigma\,\mathbf{b}'(0) + \frac{\sigma^2}{2}\,\mathbf{b}''(0) + \frac{\sigma^3}{6}\,\mathbf{b}'''(0) + ...,\end{aligned}$$

where

$$\mathbf{a}'(0)\cdot\mathbf{a}''(0) = \mathbf{b}'(0)\cdot\mathbf{b}''(0) = 0$$

due to the normality conditions $\|\mathbf{a}'(\sigma)\| = \|\mathbf{b}'(\sigma)\| = 1$.

At the second order in σ, a loop is formed when

$$\mathbf{a}(\sigma_1 - t) + \mathbf{b}(\sigma_1 + t) = \mathbf{a}(\sigma_2 - t) + \mathbf{b}(\sigma_2 + t)$$

or

$$(\sigma_1 + \sigma_2 - 2t)\,\mathbf{a}''(0) = (\sigma_1 + \sigma_2 + 2t)\,\mathbf{b}''(0),$$

which is satisfied only if $\sigma_1 = \sigma_2 = t$. Thus, the small loops could not be formed for $t \neq 0$ at the scales at which the third derivative of $\mathbf{a}(\sigma)$ or $\mathbf{b}(\sigma)$ is negligible.

The relevant statistical quantities are given by

$$\begin{aligned}\lambda_2 &\equiv \langle \mathbf{A}'(\sigma)\rangle_{rms}\\ &= \langle \int_{\sigma}^{\sigma+\Delta} \mathbf{a}''(x)dx\rangle_{rms} =\\ &= \langle \mathbf{a}'(\sigma+\Delta)-\mathbf{a}'(\sigma)\rangle_{rms}\end{aligned}$$

and

$$\begin{aligned}\lambda_3 &\equiv \langle \mathbf{A}''(\sigma)\rangle_{rms}\\ &= \langle \int_{\sigma}^{\sigma+\Delta} \mathbf{a}'''(x)dx\rangle_{rms} =\\ &= \langle \mathbf{a}''(\sigma+\Delta)-\mathbf{a}''(\sigma)\rangle_{rms}.\end{aligned}$$

The production of loops of size Δ in the vicinity of a cusp is allowed only if the following condition is satisfied

$$\lambda_3(\Delta)\frac{\Delta^3}{6} \sim \lambda_2(\Delta)\frac{\Delta^2}{2}$$

There is a divergence at small scales in the total number of loops if

$$\lim_{\Delta\to 0}\frac{\lambda_3\Delta}{3\lambda_2} > 0,$$

but the total energy emitted by a single cusp remains always finite. In the absence of divergences the smallest size of loops α_s could be approximately obtained from

$$\frac{\lambda_3(\alpha_s)\alpha_s}{3\lambda_2(\alpha_s)} \approx 0.1, \tag{2}$$

which correspond to the smallest scale at which the third term in the expansion of $\mathbf{a}(\sigma)$ remains of the same order as the second term.

IV. SPECTRUM OF LOOPS

Without loss of generality, in the reminder of the paper we calculate the spectrum of loops at a fixed time $t = 1$, assuming that the full scaling was already established. This is done to remove a trivial dependence of the scaling production function of loops on time $f(\frac{\Delta}{t}) \to f(\Delta)$.

In between the small (α_s) and large (α_l) cut-off scales there are three different regimes in the production spectrum of loops, which correspond to the populations of small, medium and large loops. The production function of loops could be estimated as

$$f(\Delta)\Delta^2 \propto \frac{\lambda_2(\Delta)}{V(\Delta)}\Delta^2, \tag{3}$$

as it must be inversely proportional to the volume V, where the vectors $\mathbf{A}(\sigma)$ and $-\mathbf{B}(\sigma)$ live, and directly proportional to the rate λ_2 at which the volume V is explored by these vectors. The two factors of Δ are due to the fact that we are interested in the energy emitted into logarithmic bins of the loop length.

From Eq. 3 we obtain

$$\begin{aligned}f_{small}(\Delta)\Delta^2 &\propto \frac{\lambda_2(\Delta)}{V_{small}(\Delta)}\Delta^2 = \frac{\lambda_2(\Delta)}{4\pi\delta(\Delta)}\\ f_{medium}(\Delta)\Delta^2 &\propto \frac{\lambda_2(\Delta)}{V_{medium}(\Delta)}\Delta^2 = \frac{3\lambda_2(\Delta)}{4\pi\Delta}\\ f_{large}(\Delta)\Delta^2 &\propto \frac{\lambda_2(\Delta)}{V_{large}(\Delta)}\Delta^2 = \frac{3\lambda_2(\Delta)}{4\pi\xi^{\frac{3}{2}}}\sqrt{\Delta}.\end{aligned}$$

The numerical analysis of Ref. [18], showed little dependence of the intercommutation probability p on the correlation length ξ, which is a hidden effect of the cross-correlations to be discussed in the last section. Deep in the scaling regime, when the vectors $\mathbf{A}'(\sigma)$ are correlated over large distances, the structures undergo a multiple intersections. This leads to a very high effective probability of their intercommutation regardless of p. On the other hand, the nearby strings inter-commute with very low probabilities, which causes the interstring distance to vary with p. Let us assume that

$$d \propto \xi\, p^{\kappa},$$

where κ is yet unknown parameter.

In the scaling regime the total energy density in long strings scales as

$$\rho \equiv \frac{1}{d^2} \propto \frac{1}{\xi^2 p^{2\kappa}},$$

but the effective intercommutation probability remains exactly one. Therefore, the probability of a single intercommutation must be proportional to the density of long strings, which implies that the production functions of loops is

$$f(\Delta)\Delta^2 \propto p^{-2\kappa}\frac{\lambda_2(\Delta)}{V(\Delta)}\Delta^2, \tag{4}$$

where p can be arbitrary small.

For very low p, the loops of much large sizes are allowed to form without ever rejoining back to the infinite network. Thus, the large cut-off scale, α_l, can be sensitive to the intercommutation probability. A very large Brownian loop of size Δ would sweep an area with radius $r \sim \xi\sqrt{\frac{\Delta}{\xi}}$ with about $\frac{r^2}{d^2}$ strings passing through the area. Each of these strings would intersect the Brownian loop roughly $\frac{r}{d}$ times, and thus,

$$\alpha_l \sim \xi\, p^{\frac{2}{3}(3\kappa-1)}. \tag{5}$$

It follows that the total energy emitted into large loops is given by

$$E_{total} \propto p^{-2\kappa}\int_0^{\xi\, p^{\frac{2}{3}(3\kappa-1)}} \frac{\lambda_2(\Delta)}{V_{large}(\Delta)}\Delta^2 d\ln(\Delta),$$

At the same time, if the decay is dominated by production of large loops, then $E_{total} \propto p^{-2\kappa}$. Therefore, the above integral must not depend on p, which implies that $\kappa = \frac{1}{3}$ and

$$d \propto \xi p^{\frac{1}{3}}.$$

Remarkably, this result is in a very good agreement with numerical simulation of Ref. [18], where $\kappa \approx 0.31$.

The above analysis shows that the cut-off scale of large loops, α_l, does not depend on p, and is always given by Eq. 1. The cut-off scale of small loops, α_s, is determined from the statistical properties of strings which was shown not to depend on p. Since the production functions of loops scale as $p^{-2\kappa} = p^{-2/3}$ and the cut-off scales are independent of p, the over all spectrum of loops must also scale as $p^{-2/3}$.

V. EXPONENTIAL APPROXIMATION

In what follows, we assume that the two-point correlation function $C(\sigma)$ is approximated by

$$C(\sigma) = e^{-a|\sigma|}.$$

The first two parameters which describe the statistics of segments $\mathbf{A}(\sigma)$ and $-\mathbf{B}(\sigma)$ are calculated exactly

$$\begin{aligned}\lambda_1(\Delta) &= \sqrt{\int_0^\Delta e^{-a|\sigma|}dx\,dy} \\ &= \sqrt{\frac{2\Delta}{a} - \frac{2}{a^2}(e^{-a\Delta} - 1)} \\ \lambda_2(\Delta) &= \sqrt{2 - 2C(\Delta)} = \sqrt{a\Delta} \\ \lambda_3(\Delta) &= \sqrt{2a^2(1 - e^{-a\Delta})}\end{aligned}$$

and the third parameter is approximated in the limit of small loops

$$\delta(\Delta) = \sqrt{\lambda_1^2 - (\int_0^\Delta C(\sigma))^2} = \sqrt{\frac{2a\Delta^3}{3}}.$$

Without the cross-correlations between opposite-moving waves taking into account, one can make an estimate of the production function of loops in the three regimes

$$\begin{aligned}f_{small}(\Delta)\Delta^2 &\propto \frac{p^{-2\kappa}}{4\pi}\sqrt{\frac{3}{2}}\Delta^{-1} \\ f_{medium}(\Delta)\Delta^2 &\propto \frac{3p^{-2\kappa}}{4\pi}\sqrt{a}\Delta^{-\frac{1}{2}} \\ f_{large}(\Delta)\Delta^2 &\propto \frac{3p^{-2\kappa}}{4\pi}\sqrt{a}\xi^{-\frac{3}{2}}\Delta\end{aligned} \tag{6}$$

The energy scale of small loops can be estimated from Eq. 2:

$$\alpha_s \approx \frac{0.15}{a}.$$

There are two more relevant scales α_{m_1} and α_{m_2}, which specify the range of the medium loops. From Eqs. 6 $\alpha_{m_1} = (6\,a)^{-1}$ and $\alpha_{m_2} = \xi$.

An important ratio is given by

$$\beta \equiv \frac{\int_{\alpha_s}^{\alpha_{m_1}} f_{small}(\Delta)\Delta^2 d\ln(\Delta)}{E_{total}}$$

where

$$\begin{aligned}E_{total} \;=\; &\int_{\alpha_s}^{\alpha_{m_1}} f_{small}(\Delta)\Delta^2 d\ln(\Delta) + \\ &\int_{\alpha_{m_1}}^{\alpha_{m_2}} f_{medium}(\Delta)\Delta^2 d\ln(\Delta) + \\ &\int_{\alpha_{m_2}}^{\alpha_l} f_{large}(\Delta)\Delta^2 d\ln(\Delta),\end{aligned}$$

which roughly determines the fraction of energy emitted into small loops. For $\xi = 0.1$, $a = 3$ we obtain

$$\begin{aligned}\alpha_l &\sim 0.3, \\ \alpha_s &\sim 0.05\end{aligned}$$

and

$$\beta \sim 0.5.$$

Thus, about 50% of the energy goes into small loops in contrast to 90% obtained in Ref. [19], where the power-law approximation of the two point correlation was assumed. Our model with the power-law ansatz also leads to immediate divergences at small scales. In a real network λ_3 can in principle be divergent when the kinks are present. However, our analysis assumes the Taylor expansion only around a given cusp. The probability of a near-cusp region to contain recent kinks arbitrary close to the cusp is infinitely small. [1] We conclude that physically relevant λ_3 cannot be divergent, and $C(\sigma)$ should not be modeled by a power-law.

The spectrum of loops for the above example is shown on Fig. 1. The qualitative shape of the plot agrees very well with numerical simulation of Refs. [11, 12]. Nevertheless, it is too early to expect a full quantitative agreement between our analytic model and different simulations, due to a couple of reasons. First of all, at present time all of the simulations are not entirely in the scaling regime, as the production of loops is dominated by the remnants of the initial conditions. Secondly, our model, as it stands, is ignorant to the cross-correlations between opposite-moving waves. In the following section, we will make a first attempt to include such effects into our formalism.

---

[1] The old kinks are completely smoothed out by gravitational radiation after a fixed (may be very large) time, but the frequency of cusps formation decreases with time. Therefore, in the long run, the old kinks can be completely ignored for the analysis of small loops formation in the vicinity of cusps.



Figure 1: Production function of loops.

VI. DISCUSSION

In the limit of negligible expansion the left and right moving waves stay practically unchanged throughout the evolution. Therefore,

$$\mathbf{A}(0) - \mathbf{B}(\sigma) > \epsilon$$

must be satisfied for all $0 < \sigma < \xi$, or otherwise a loop of size Δ would have been formed already. Given that, we want to find the probability P that the following inequality holds:

$$\mathbf{A}(0) - \mathbf{B}(0) < \epsilon.$$

At the moment, let us only consider the small loops. We expect P to be smaller, than one would get without cross-correlations taken into account, as it was done in the previous sections. On the other hand, if another intercommutation is taking place such that

$$\mathbf{A}(\sigma) - \mathbf{B}(\sigma) < \epsilon$$

for some $-\xi < \sigma < 0$, than P must be much larger. The over all effect on the production of small loops will be the same as if no cross-correlations are present.

Although, the production of small loops is a very rare process as the correlation length grows, the total effect is compensated by a large number of loops produced at once. It follows that a number of small loops are likely to be emitted in bursts, with very high velocities pointing in approximately the same direction. These loops are not likely to fragment further, since the third derivative of the segments at even smaller scales ($< \alpha_s$) is not sufficiently large, and as a result the small loops will decay only gravitationally.

For the large and medium loops the situation is more complicated and at present it is not clear how to take the effect of cross-correlations into account. In addition, it is desired to include the intersections of nearby strings, which inter-commute and produce kinks. All of these effects complicate the problem dramatically and can in principle lead to some unexpected results. At present we simply assume that the production of large and medium-sized loops is captured entirely by the statistics of long strings, in analogy to the small loops.

To conclude, we summarize the main differences of our results from those obtained in Refs. [16, 17]. In the scaling regime, we do not expect to see very small loops at the scale of the gravitational back reaction. Most of the energy is emitted into two different populations of the string loops: small and large. The scale of large loops is set by the correlation length, which does not depend on the intercommutation probability. The scale of small loops is determined from the statistics of structures on infinite strings, and is expected to be only few orders lower than the scale of large loops. For illustration we used an exponential form of the correlation function in contrast to the power-law approximation used in Refs. [16, 17], which cannot correctly represent a physically relevant statistics of strings at small scales.

Acknowledgments

We are grateful to Alex Vilenkin, Ken Olum and Joe Polchinski for very helpful discussions and comments on the manuscript. This work was supported in part by the Transregional Collaborative Research Centre TRR 33 "The Dark Universe ".

[1] T. W. B. Kibble, J. Phys. **A9**, 1387 (1976); For a review see A. Vilenkin and E. P. S. Shellard, *Cosmic Strings and Other Topological Defects* (Cambridge University Press, 2000).
[2] S. Sarangi and S. H. Tye, Phys. Lett. **B536**, 185 (2002).
[3] G. Dvali and A. Vilenkin, JCAP **0403**, 010 (2004).
[4] E. J. Copeland, R. C. Myers and J. Polchinski, JHEP **06**, 013 (2004).
[5] T. W. B. Kibble, Nucl. Phys. B **252**, 227 (1985) [Erratum-ibid. B **261**, 750 (1985)].
[6] D. P. Bennett, Phys. Rev. D **34**, 3592 (1986).
[7] A. Albrecht and N. Turok, Phys. Rev. Lett. **54**, 1868 (1985); Phys. Rev. D **40**, 973 (1989).
[8] D. P. Bennett and F. R. Bouchet, Phys. Rev. Lett. **60**, 257 (1988); **63**, 2776 (1989); Phys. Rev. D **41**, 2408 (1990).
[9] B. Allen and E. P. S. Shellard, Phys. Rev. Lett. **64**, 119 (1990)

[10] V. Vanchurin, K. Olum and A. Vilenkin, Phys. Rev. D **72**, 063514 (2005)

[11] V. Vanchurin, K. D. Olum and A. Vilenkin, Phys. Rev. D **74**, 063527 (2006)

[12] K. D. Olum and V. Vanchurin, Phys. Rev. D **7**5, 063521 (2007)

[13] C. J. A. P. Martins and E. P. S. Shellard, Phys. Rev. D **73**, 043515 (2006)

[14] C. Ringeval, M. Sakellariadou and F. Bouchet, JCAP 0702:023 (2007)

[15] T. Vachaspati and A. Vilenkin, Phys. Rev. D **30**, 2036 (1984)

[16] F. Dubath, J. Polchinski and J. V. Rocha, arXiv:0711.0994

[17] J. Polchinski and J. V. Rocha, Phys. Rev. D75, 123503 (2007)

[18] V. Vanchurin, Ph.D. Thesis: “Evolution of cosmic strings,” UMI-31-82849 (2005)

[19] J. Polchinski, arXiv:0707.0888